

## PERSIMMON

### PROCEDURES FOR SOIL AND LEAF SAMPLING

Soil and leaf analyses can help determine the fertilization needs of your crops and assist in identifying problem conditions if they exist. This guide will help you collect representative and meaningful samples which will assure accurate laboratory analysis data.

Leaf analysis is most useful in evaluating the nutrient status or toxic levels of a growing crop especially when results can be compared with interpretive guides.

Preplant or postplant soil analyses provide useful information for measurement of fertility or toxicity levels and to substantiate amendment requirements. Data from soil analyses is most meaningful when used in conjunction with leaf tissue analysis results.

### SOIL SAMPLING - PERSIMMON

#### When to Sample - Soils

Preplant soil analysis data is used to determine the most appropriate root stock and helps determine the need for preplant fertilizers or soil amendments.

Soil samples in established plantings are generally collected in spring. However, soil samples may be collected throughout the year to monitor soil conditions or to determine the causes for poor growth in specific locations.

#### How to Sample - Soils

Soil samples should be representative of the area to be treated. If possible, areas should be uniform with respect to soil texture, slope, variety and irrigation design. Areas sampled should not be larger than 10-20 acres. Problem areas should be sampled separately and compared with samples taken from adjacent non problem areas. The location of the sample areas should be noted and marked on a parcel or planting map for future reference.

Soil samples should contain at least 10-20 cores for each designated area. Samples should be collected from the irrigated area, at or near the drip line of the trees. Retrieve an equal amount of soil from the surface to a depth of 24 inches (0-24"). Soil core samples should be placed in a clean bucket and thoroughly mixed. Approximately a quart volume of this soil will be required for analysis purposes.



## **LEAF SAMPLING - PERSIMMON**

### **When to Sample - Leaves**

Persimmon leaf samples are collected in July to September. For problem diagnosis other sampling times are acceptable.

### **How to Sample - Leaves**

Leaf samples should be representative of the area to be fertilized. The sampler should traverse each sample area collecting samples perpendicularly or diagonally to the tree rows. Sampling areas should be uniform with respect to soil texture, slope, variety and irrigation design and should not be larger than 10-20 acres. Problem areas should be sampled separately.

### **Volume and Type of Leaves to Collect**

25 leaves from each area will be sufficient. Collect fully expanded mature leaves (two to five months old) from non-fruiting spurs.

### **Labeling, Packaging, and Shipping**

All leaf and soil samples should be labeled with your name and address, sample identification, crop, tree age, previous problems (if any) and the required analysis. Leaf tissues should be placed in paper bags and soil samples in plastic bags. Leaf samples should be delivered to the laboratory as soon as possible to insure freshness. If held for more than one day, leaf samples should be stored in paper bags at room temperature until delivered or shipped. Analytical nutrient values are not affected if samples are allowed to air dry naturally. This becomes a problem when moisture is present and mold develops.

As an alternative to delivering directly to our laboratory, leaf and soil samples can be shipped via overnight UPS or other private couriers.

### **When in doubt.....**

If you have any questions or require assistance relating to the above, please visit [www.fglinc.com](http://www.fglinc.com) or call Fruit Growers Laboratory's Agronomic Services.